

# ToO update from MIRO



**Jet Propulsion Laboratory**  
California Institute of Technology

Anthony Lethuillier<sup>1</sup>, Paul von Allmen<sup>1</sup>, Mark Hofstadter<sup>1</sup>, Gerard Beaudin<sup>4</sup>, Nicolas Biver<sup>2</sup>, Dominique Bockelee-Morvan<sup>2</sup>, Mathieu Choukroun<sup>1</sup>, Jacques Crovisier<sup>2</sup>, Bjorn Davidsson<sup>1</sup>, Pierre Encrenaz<sup>4</sup>, Therese Encrenaz<sup>2</sup>, Margaret Frerking<sup>1</sup>, Adeline Gicquel<sup>1</sup>, Samuel Gulkis<sup>1</sup>, Paul Hartogh<sup>5</sup>, Wing-Huen Ip<sup>6</sup>, Michael A. Janssen<sup>1</sup>, Christopher Jarchow<sup>5</sup>, Seungwon Lee<sup>1</sup>, Emmanuel Lellouch<sup>2</sup>, Cedric Leyrat<sup>2</sup>, Ladislav Rezac<sup>5</sup>, Peter Schloerb<sup>3</sup>, Thomas R. Spilker<sup>7</sup>

<sup>1</sup>Jet Propulsion Laboratory/Calif. Inst. Tech., Pasadena, CA, United States. <sup>2</sup>LESIA-Observatoire de Paris, Meudon, 92195, France. <sup>3</sup>University of Massachusetts, Amherst, MA, United States. <sup>4</sup>LERMA-Observatoire de Paris, Paris, 75014, France. <sup>5</sup>Max-Planck-Institut für Sonnensystemforschung, Göttingen, 37077, Germany. <sup>6</sup>National Central University, Taoyuan City, 32001, Taiwan. <sup>7</sup>Solar System Science and Exploration, Los Angeles, CA, United States.

© 2017 California Institute of Technology. Government sponsorship acknowledged.



## Context and Model

---

# Context

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

## The first ToO:

- Is Located in the Imhotep region of the comet.




Figure 1: NAVCAM image of the Imhotep region with indicated the 2014 and 2016 swaths. Copyright: ESA/Rosetta/NAVCAM

# Context

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

## The first ToO:

- Is Located in the Imhotep region of the comet.
- First observation: October 27th 2014 as a single swath.




Figure 1: NAVCAM image of the Imhotep region with indicated the 2014 and 2016 swaths. Copyright: ESA/Rosetta/NAVCAM

# Context

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

## The first ToO:

- Is Located in the Imhotep region of the comet.
- First observation: October 27th 2014 as a single swath.
- Second observation: July 9th 2016 as a raster scan.




Figure 1: NAVCAM image of the Imhotep region with indicated the 2014 and 2016 swaths. Copyright: ESA/Rosetta/NAVCAM

# Context

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

## The first ToO:

- Is Located in the Imhotep region of the comet.
- First observation: October 27th 2014 as a single swath.
- Second observation: July 9th 2016 as a raster scan.
- Objective: observe the same area twice with similar high spatial resolution.





Figure 1: NAVCAM image of the Imhotep region with indicated the 2014 and 2016 swaths. Copyright: ESA/Rosetta/NAVCAM

# Measurements

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives



Resolution	SMM	MM
October 2014	$\approx 20m$	$\approx 60m$
July 2016	$\approx 30m$	$\approx 90m$

The 2016 raster scan intersected the 2014 swath a total of 14 times.

*Figure 2: NAVCAM image of the Imhotep region with indicated the 2014 and 2016 swaths. Copyright: ESA/Rosetta/NAVCAM*

# Thermal and radiative model

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives




Figure 3: Simplified thermal model of the subsurface of 67P/C-G, indicating the processes at play



# Thermal and radiative model

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

Parameters	MM	SMM
Pure dust density ( $\text{Kg}/\text{m}^3$ )	2500	
Pure ice density ( $\text{Kg}/\text{m}^3$ )	925	
Emissivity	0.9	
Albedo	0.04	
Penetration depth	$\approx 4\text{cm}$	$\approx 1\text{cm}$
Refraction index	$3.08 - j0.051$	$3.08 - j0.017$
Upper density	Variable	
Lower density		
Dust/ice ratio		




## Results and Interpretations

---



# Best fitting parameters in 2014 and 2016

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives



# Best fit for both times

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives



Anthony Lethuillier 06/11/2017 SWT Rosetta

# Thermal inertia of the subsurface

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

Instrument	Region (date)	Thermal Inertia
MIRO <sup>1</sup>	All nucleus (2014)	10 – 50
MIRO <sup>2</sup>	Imhotep and Ash (September 2014)	10 – 30
MUPUS <sup>3</sup>	Abydos (November 2014)	50 – 120
MIRO <sup>4</sup>	Imhotep ToO (September 2014)	80 – 95

<sup>1</sup> Gulkis et al. (2015) <sup>2</sup> Schloerb et al. (2015) <sup>3</sup> Spohn et al. (2015) <sup>4</sup> This work



# Porosity of the subsurface

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

Instrument	Depth sounded	Porosity
CONSERT <sup>1</sup>	Hundreds of meters	75 – 85%
SESAME-PP <sup>2</sup>	First meter	< 50%
MIRO <sup>4</sup>	10 cm	50 – 60%
MUPUS <sup>3</sup>	Near surface	30 – 65%

<sup>1</sup> Kofman et al. (2015) <sup>2</sup> Lethuillier et al. (2016) <sup>3</sup> Spohn et al. (2015) <sup>4</sup> This work





## **Conclusions and Perspectives**

---

# Conclusion and perspectives for the Imhotep ToO

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

- The model used in this work offers a powerful tool to investigate the subsurface of 67P/C-G using MIRO data and can help constrain the composition of the subsurface and its evolution through time.





# Conclusion and perspectives for the Imhotep ToO

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

- The model used in this work offers a powerful tool to investigate the subsurface of 67P/C-G using MIRO data and can help constrain the composition of the subsurface and its evolution through time.
- More parameters need to be investigated in order to better fit the measurements of the Imhotep ToO.



# Conclusion and perspectives for the Imhotep ToO

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

- The model used in this work offers a powerful tool to investigate the subsurface of 67P/C-G using MIRO data and can help constrain the composition of the subsurface and its evolution through time.
- More parameters need to be investigated in order to better fit the measurements of the Imhotep ToO.
- For this a MCMC algorithm will be used in future work.



# Conclusion and perspectives for the Imhotep ToO

1. Context and Model
2. Results and Interpretations
3. Conclusions and Perspectives

- The model used in this work offers a powerful tool to investigate the subsurface of 67P/C-G using MIRO data and can help constrain the composition of the subsurface and its evolution through time.
- More parameters need to be investigated in order to better fit the measurements of the Imhotep ToO.
- For this a MCMC algorithm will be used in future work.
- MIRO made observations of Imhotep at other times, and these will be included in future studies.

